


Application No. 10/772,069

AMENDMENTS TO THE CLAIMS

A detailed listing of all claims that are, or were, in the present application, irrespective of whether the claim(s) remains under examination in the application are presented below. The claims are presented in ascending order and each includes one status identifier.

1. (Withdrawn) An organophotoreceptor comprising:

- (a) a charge transport composition comprising molecules having the formula



where n is an average of a distribution of integers in which n is at least 2;

R₁, R₂, R₃, and R₄ comprise, each independently, H, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group;

X comprises an (N,N-disubstituted)arylamine group; and

Y is a bridging group;

- (b) a charge generating compound; and


- (c) an electrically conductive substrate over which the charge transport composition and the charge generating compound are located.

2. (Withdrawn) An organophotoreceptor according to claim 1 wherein X is selected from the group consisting of a carbazole group, a julolidine group, a triarylamine group, a dialkylarylamine group, and an alkyl diarylamine group.

Application No. 10/772,069

3. (Withdrawn) An organophotoreceptor according to claim 1 wherein Y comprises a $-(CH_2)_m-$ group where m is an integer between 1 and 30, inclusive, and one or more of the methylene groups is optionally replaced by O, S, N, C, Si, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, urethane, urea, an ester group, an NR_5 group, a CR_6 , or a CR_7R_8 group where R_5 , R_6 , R_7 , and R_8 are, each independently, a bond, H, hydroxyl, thiol, carboxyl, an amino group, an alkyl group, an alkenyl group, a heterocyclic group, an aromatic group, or part of a ring group.

4. (Withdrawn) An organophotoreceptor according to claim 3 wherein Y is selected from the group consisting of the formulae:



where Q, X_1 , X_2 , X_3 , and X_4 are, each independently, O, S, or NR' where R' comprises H, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group.

5. (Withdrawn) An organophotoreceptor according to claim 1 further comprising an electron transport compound.

6. (Withdrawn) An organophotoreceptor according to claim 1 wherein said organophotoreceptor is in the form of a drum or a belt.

Application No. 10/772,069

7. (Withdrawn) An organophotoreceptor according to claim 1 comprising:

- (a) a charge transport layer comprising said charge transport composition and a polymeric binder; and
- (b) a charge generating layer comprising said charge generating compound and a polymeric binder.


8. (Withdrawn) An organophotoreceptor according to claim 1 wherein n is at least

5.

9. (Withdrawn) An electrophotographic imaging apparatus comprising:

- (a) a light imaging component; and
- (b) an organophotoreceptor oriented to receive light from the light imaging component, the organophotoreceptor comprising an electrically conductive substrate and a photoconductive element on the electrically conductive substrate, the photoconductive element comprising:

(i) a charge transport composition comprising molecules having the formula



where n is an average of a distribution of integers in which n is at least 2;

R₁, R₂, R₃, and R₄ comprise, each independently, H, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group;

X comprises an (N,N-disubstituted)arylamine group; and

Y is a bridging group; and


(ii) a charge generating compound.

Application No. 10/772,069

10. (Withdrawn) An electrophotographic imaging apparatus according to claim 9 wherein X is selected from the group consisting of a carbazole group, a julolidine group, a triarylamine group, a dialkylarylamine group, and an alkyl diarylamine group.

11. (Withdrawn) An electrophotographic imaging apparatus according to claim 9 wherein Y comprises a $-(CH_2)_m-$ group where m is an integer between 1 and 30, inclusive, and one or more of the methylene groups is optionally replaced by O, S, N, C, Si, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, urethane, urea, an ester group, an NR_5 group, a CR_6 , or a CR_7R_8 group where R_5 , R_6 , R_7 , and R_8 are, each independently, a bond, H, hydroxyl, thiol, carboxyl, an amino group, an alkyl group, an alkenyl group, a heterocyclic group, an aromatic group, or part of a ring group.

12. (Withdrawn) An electrophotographic imaging apparatus according to claim 11 wherein Y is selected from the group consisting of the formulae:



where Q, X₁, X₂, X₃, and X₄ are, each independently, O, S, or NR' where R' comprises H, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group.

Application No. 10/772,069

13. (Withdrawn) An electrophotographic imaging apparatus according to claim 9 comprising a toner dispenser.


14. (Withdrawn) An electrophotographic imaging apparatus according to claim 9 further comprising an electron transport compound.

15. (Withdrawn) An electrophotographic imaging apparatus according to claim 9 wherein n is at least 5.

16. (Withdrawn) An electrophotographic imaging process comprising:

(a) applying an electrical charge to a surface of an organophotoreceptor comprising an electrically conductive substrate and a photoconductive element on the electrically conductive substrate, the photoconductive element comprising:

(i) a charge transport composition comprising molecules having the formula



where n is an average of a distribution of integers in which n is at least 2;

R₁, R₂, R₃, and R₄ comprise, each independently, H, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group;

X comprises an (N,N-disubstituted)arylamine group; and

Y is a bridging group; and

(ii) a charge generating compound;

Application No. 10/772,069

(b) imagewise exposing the surface of the organophotoreceptor to radiation to dissipate charge in selected areas and thereby form a pattern of charged and uncharged areas on the surface;


(c) contacting the surface with a toner to create a toned image; and

(d) transferring the toned image to a substrate.


17. (Withdrawn) An electrophotographic imaging process according to claim 16 wherein X is selected from the group consisting of a carbazole group, a julolidine group, a triarylamine group, a dialkylarylamine group, and an alkyl diarylamine group.

18. (Withdrawn) An electrophotographic imaging process according to claim 16 wherein Y comprises a $-(CH_2)_m-$ group where m is an integer between 1 and 30, inclusive, and one or more of the methylene groups is optionally replaced by O, S, N, C, Si, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, urethane, urea, an ester group, an NR_5 group, a CR_6 , or a CR_7R_8 group where R_5 , R_6 , R_7 , and R_8 are, each independently, a bond, H, hydroxyl, thiol, carboxyl, an amino group, an alkyl group, an alkenyl group, a heterocyclic group, an aromatic group, or part of a ring group.

19. (Withdrawn) An electrophotographic imaging process according to claim 18 wherein Y is selected from the group consisting of the formulae:



Application No. 10/772,069




where Q, X₁, X₂, X₃, and X₄ are, each independently, O, S, or NR' where R' comprises H, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group.

20. (Withdrawn) An electrophotographic imaging process according to claim 16 wherein the toner comprises a dispersion of colorant particles.

21. (Withdrawn) An electrophotographic imaging process according to claim 16 further comprising an electron transport compound.

22. (Withdrawn) An electrophotographic imaging process according to claim 14 wherein n is at least 5.

23. (Original) A charge transport composition comprising molecules having the formula



where n is an average of a distribution of integers in which n is at least 2;

R₁, R₂, R₃, and R₄ comprise, each independently, H, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group;

X comprises an (N,N-disubstituted)arylamine group; and


Y is a bridging group.

Application No. 10/772,069

24. (Original) A charge transport composition according to claim 23 wherein X is selected from the group consisting of a carbazole group, a julolidine group, a triarylamine group, a dialkylarylamine group, and an alkyl diarylamine group.

25. (Original) A charge transport composition according to claim 23 wherein Y comprises a $-(CH_2)_m-$ group where m is an integer between 1 and 30, inclusive, and one or more of the methylene groups is optionally replaced by O, S, N, C, Si, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, urethane, urea, an ester group, an NR₅ group, a CR₆, or a CR₇R₈ group where R₅, R₆, R₇, and R₈ are, each independently, a bond, H, hydroxyl, thiol, carboxyl, an amino group, an alkyl group, an alkenyl group, a heterocyclic group, an aromatic group, or part of a ring group.

26. (Original) A charge transport composition according to claim 25 wherein Y is selected from the group consisting of the formulae:




where Q, X₁, X₂, X₃, and X₄ are, each independently, O, S, or NR' where R' comprises H, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group.

Application No. 10/772,069

27. (Original) A charge transport composition according to claim 25 wherein n is at least 5.

28. (Original) A charge transport composition prepared by the reaction of a multi-functional compound with a di-reactive-ring compound having the formula



where R_1' , R_2' , R_3' , and R_4' comprise, each independently, H, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group;

X' comprises an (N,N-disubstituted)arylamine group; and

Y_1 and Y_2 are, each independently, a linking group; and

E_1 and E_2 are, each independently, a reactive ring group.

29. (Original) A charge transport composition according to claim 28 wherein X' is selected from the group consisting of a carbazole group, a julolidine group, a triarylamine group, a dialkylarylamine group, and an alkyldiarylamine group.

30. (Original) A charge transport composition according to claim 28 wherein Y_1 and Y_2 , each independently, comprise a $-(CH_2)_k-$ group where k is an integer between 1 and 30, inclusive, and one or more of the methylene groups is optionally replaced by O, S, N, C, Si, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, urethane, urea, an ester group, an NR_{16} group, a CR_{17} , or a $CR_{18}R_{19}$ group where R_{16} , R_{17} , R_{18} , and R_{19} are, each independently, a bond,

Application No. 10/772,069

H, hydroxyl, thiol, carboxyl, an amino group, an alkyl group, an alkenyl group, a heterocyclic group, an aromatic group, or part of a ring group.

31. (Original) A charge transport composition according to claim 28 wherein E_1 and E_2 , each independently, are selected from the group consisting of 3-, 4-, 5-, 7-, 8-, 9-, 10-, 11- and 12-membered heterocyclic ring groups.

32. (Original) A charge transport composition according to claim 31 wherein E_1 and E_2 , each independently, are selected from the group consisting of 3-, 4-, 5-, 7-, 8-, 9-, 10-, 11-, and 12-membered cyclic ethers, cyclic amines, cyclic sulfides, cyclic amides, N-carboxy-amino acid anhydrides, lactones, and cyclosiloxanes.

33. (Original) A charge transport composition according to claim 32 wherein E_1 and E_2 , each independently, are selected from the group consisting of epoxides, oxetanes, aziridines, thiiranes, 2-azetidinone, 2-pyrrolidone, 2-piperidone, caprolactam, enantholactam, and capryllactam.


34. (Original) A charge transport composition according to claim 28 wherein the multi-functional compound is a di-functional compound.

35. (Original) A charge transport composition according to claim 34 wherein the di-functional compound is selected from the group consisting of a triol, a triamine, a trithiol, a diol, a dithiol, a diamine, a dicarboxylic acid, a hydroxylamine, an amino acid, a hydroxyl acid, a thiol acid, a hydroxythiol, and a thioamine.

Application No. 10/772,069

36. (Withdrawn) An organophotoreceptor comprising:

(a) a polymeric charge transport composition prepared by the reaction of a multi-functional compound with a di-reactive-ring compound having the formula



where R_1' , R_2' , R_3' , and R_4' comprise, each independently, H, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group;

X' comprises an (N,N-disubstituted)arylamine group; and

Y_1 and Y_2 are, each independently, a linking group; and

E_1 and E_2 are, each independently, a reactive ring group; and

(b) a charge generating compound; and

(c) an electrically conductive substrate over which the charge transport composition and the charge generating compound are located.

37. (Withdrawn) An organophotoreceptor according to claim 36 wherein X' is selected from the group consisting of a carbazole group, a julolidine group, a triarylamine group, a dialkylarylamine group, and an alkyldiarylamine group.

38. (Withdrawn) A charge transport composition according to claim 36 wherein Y_1 and Y_2 , each independently, comprise a $-(CH_2)_k-$ group where k is an integer between 1 and 30, inclusive, and one or more of the methylene groups is optionally replaced by O, S, N, C, Si, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, urethane, urea, an ester group, an NR_{16} group, a CR_{17} , or a $CR_{18}R_{19}$ group where R_{16} , R_{17} , R_{18} , and R_{19} are, each independently, a bond,

Application No. 10/772,069

H, hydroxyl, thiol, carboxyl, an amino group, an alkyl group, an alkenyl group, a heterocyclic group, an aromatic group, or part of a ring group.

39. (Withdrawn) A charge transport composition according to claim 36 wherein E_1 and E_2 , each independently, are selected from the group consisting of 3-, 4-, 5-, 7-, 8-, 9-, 10-, 11- and 12-membered heterocyclic ring groups.

40. (Withdrawn) A charge transport composition according to claim 39 wherein E_1 and E_2 , each independently, are selected from the group consisting of 3-, 4-, 5-, 7-, 8-, 9-, 10-, 11- and 12-membered cyclic ethers, cyclic amines, cyclic sulfides, cyclic amides, N-carboxy-amino acid anhydrides, lactones, and cyclosiloxanes.

41. (Withdrawn) A charge transport composition according to claim 40 wherein E_1 and E_2 , each independently, are selected from the group consisting of epoxides, oxetanes, aziridines, thiranes, 2-azetidinone, 2-pyrrolidone, 2-piperidone, caprolactam, enantholactam, and capryllactam.

42. (Withdrawn) A charge transport composition according to claim 36 wherein the multi-functional compound is a di-functional compound.

43. (Withdrawn) A charge transport composition according to claim 36 wherein the di-functional compound is selected from the group consisting of a triol, a triamine, a trithiol, a diol, a dithiol, a diamine, a dicarboxylic acid, a hydroxylamine, an amino acid, a hydroxyl acid, a thiol acid, a hydroxythiol, and a thioamine.